




**FIGURE 1****FIGURE 2****BEST AVAILABLE COPY**

**FIGURE 3**


Solubility of linker-BSA constructs (20-22) in 10 mM ammonium bicarbonate at pH 8 (semi-log plot)





10/511893

FIGURE 4

Solubility of linker-BSA constructs (20-22) in 0.1M sodium formate at pH 4.5  
(semi-log plot)



**FIGURE 5****FIGURE 6**


10/511893

PCT/GB03/01505

WO 03/087824

5/14

FIGURE 7



BEST AVAILABLE COPY


10/511893

WO 03/087824


PCT/GB03/01505



6/14



**FIGURE 8**



**FIGURE 9**



**FIGURE 10****FIGURE 11**

**FIGURE 12****FIGURE 13**

10/511893

WO 03/087824

PCT/GB03/01505

9/14

FIGURE 14







FIGURE 15




BEST AVAILABLE COPY


**FIGURE 16****FIGURE 17****BEST AVAILABLE COPY**

**FIGURE 18**

Immulon 2HB ELISA of protein-TML:biotin-hydrazone samples probed with ExtrAvidin-HRP

**FIGURE 19**

1 2 3 4 5 6 7 8 9 10

**BEST AVAILABLE COPY**


10/511893

PCT/GB03/01505


WO 03/087824

12/14

**FIGURE 20**



**FIGURE 21**



**BEST AVAILABLE COPY**



**FIGURE 22****FIGURE 23****BEST AVAILABLE COPY**

FIGURE 24





FIGURE 25



BEST AVAILABLE COPY